


Amendment to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Currently Amended) A composition of matter comprising a compound having the following structural formula:



wherein,

R₁ and R₂ are selected from H, a hydrocarbyl having up to 20 carbon atoms, and a hydrocarbyl having up to 20 carbon atoms and substituted with a group selected from hydroxy, alkoxy, amino, substituted amino, thio, alkylthio, guanidino, ureido and heterocyclyl;

R₃ is selected from a hydrocarbyl having up to 20 carbon atoms, and a hydrocarbyl having up to 20 carbon atoms and substituted with a group selected from halo, haloalkyl, hydroxy, alkyl, alkoxy, alkylendioxy, amino, substituted amino, aminoalkyl, thio, alkylthio, guanidino, ureido, heterocyclyl, heteroaryl, and heteroarylthio;

R₄ is a substituted amino, -NR₆R₇, wherein R₆ and R₇ are selected from H and a hydrocarbyl having up to 20 carbon atoms; R₆ and R₇, with inclusion of N, may combine to form a heterocyclic ring such as indolynyl having the formula



R₅ is selected from O, S, NH, N-alkyl, N-alkenyl, N-alkynyl, N-cycloalkyl, N-aryl and N-aralkyl,
and

n is 1 to 3; and

wherein the composition is further characterized by at least one additional limitation selected from the group consisting of:

one or more methylene groups of a hydrocarbonyl group of R₃ being replaced by an oxygen atom;

R₁ being selected from alkyl and aminoalkyl;

R₁ being selected from (S)-Methyl, (R)-Methyl or (S)-Propyl;

R₁ being (S)-Aminopropyl;

R₂ being selected from (R)-Aminomethyl-(imino)-propyl, (S)-Aminomethyl-(imino)-propyl or (S)-Methylthiomethyl;

R₃ being selected from alkyl, aralkyl or substituted aralkyl;

R₃ being 3-bromophenethyl;

R₃ being 3,5 bis-(trifluoromethyl)phenethyl;

R₄ being aralkylamino;

R₄ being benzylamino;

R₆ and R₇ with inclusion of n, being heterocyclyl; and

-NR₆R₇ being isoindolyl having the formula



2. A composition according to claim 1, wherein R1, R2, R3, R6 or R7 comprises a straight chain hydrocarbonyl.

3. A composition according to claim 1, wherein R1, R2, R3, R6 or R7 comprises a branched chain hydrocarbonyl.

4. A composition according to claim 1, wherein R1, R2, R3, R6 or R7 comprises a saturated hydrocarbonyl.

5. A composition according to claim 1, wherein R1, R2, R3, R6 or R7 comprises an unsaturated hydrocarbonyl.

6. A composition according to claim 1, wherein R1, R2, R3, R6 or R7 comprises a cyclic hydrocarbonyl.

7. A composition according to claim 1, wherein R1, R2, R3, R6 or R7 comprises an acyclic hydrocarbonyl.

8. A composition according to claim 1, wherein R1, R2, R3, R6 or R7 comprises a chiral hydrocarbonyl.

9. A composition according to claim 1, wherein R1, R2, R3, R6 or R7 comprises an achiral hydrocarbonyl.

10. A composition according to claim 1, wherein R1, R2, R3, R6 or R7 comprises a substituted hydrocarbonyl.

11. (Cancelled)

12. (Cancelled)

13. (Cancelled)

14. (Cancelled)

15. (Cancelled)

16. (Cancelled)

17. (Cancelled)

18. (Cancelled)

19. (Cancelled)

20. (Cancelled)

21. (Cancelled)

22. (Cancelled)


23. (Cancelled)

24. (Cancelled)

25. (Withdrawn) The use of a composition according to any of claims 1-22 in the manufacture of preparation for administration to a human or animal subject to block or antagonize MCH receptors or to decrease food intake or to treat obesity, a metabolic disorder, an eating disorder, depression or urinary incontinence.

26. (Withdrawn) A method for a) blocking or antagonizing MCH receptors or b) decreasing food intake or c) treating obesity, a metabolic disorder, an eating disorder, depression or urinary incontinence, said method comprising the step of:

administering to the individual an effective amount of a composition that comprises a compound having the following structural formula:



wherein,

R₁ and R₂ are selected from H, a hydrocarbyl having up to 20 carbon atoms, and a hydrocarbyl having up to 20 carbon atoms and substituted with a group selected from hydroxy, alkoxy, amino, substituted amino, thio, alkylthio, guanidino, ureido and heterocyclyl;

R₃ is selected from a hydrocarbyl having up to 20 carbon atoms, and a hydrocarbyl having up to 20 carbon atoms and substituted with a group selected from halo, haloalkyl, hydroxy, alkyl, alkoxy, alkylendioxy, amino, substituted amino, aminoalkyl, thio, alkylthio, guanidino, ureido, heterocyclyl, heteroaryl, and heteroarylthio;

R₄ is a substituted amino, -NR₆R₇, wherein R₆ and R₇ are selected from H and a hydrocarbyl having up to 20 carbon atoms; R₆ and R₇, with inclusion of N, may combine to form a heterocyclic ring such as indolynyl, having the formula



R₅ is selected from O, S, NH, N-alkyl, N-alkenyl, N-alkynyl, N-cycloalkyl, N-aryl and N-aralkyl, and

n is 1 to 3.

27. (Withdrawn) A method according to claim 26, wherein R₁, R₂, R₃, R₆ or R₇ comprises a straight chain hydrocarbyl.

28. (Withdrawn) A method according to claim 26, wherein R₁, R₂, R₃, R₆ or R₇ comprises a branched chain hydrocarbyl.

29. (Withdrawn) A method according to claim 26, wherein R₁, R₂, R₃, R₆ or R₇ comprises a saturated hydrocarbyl.

30. (Withdrawn) A method according to claim 26, wherein R₁, R₂, R₃, R₆ or R₇ comprises an unsaturated hydrocarbyl.

31. (Withdrawn) A method according to claim 26, wherein R₁, R₂, R₃, R₆ or R₇ comprises a cyclic hydrocarbyl.

32. (Withdrawn) A method according to claim 26, wherein R₁, R₂, R₃, R₆ or R₇ comprises an acyclic hydrocarbyl.

33. (Withdrawn) A method according to claim 26, wherein R₁, R₂, R₃, R₆ or R₇ comprises a chiral hydrocarbyl.

34. (Withdrawn) A method according to claim 26, wherein R₁, R₂, R₃, R₆ or R₇ comprises an achiral hydrocarbyl.

35. (Withdrawn) A method according to claim 26, wherein R₁, R₂, R₃, R₆ or R₇ comprises a substituted hydrocarbyl.

36. (Withdrawn) A method according to claim 26, wherein one or more methylene groups of a hydrocarbyl group of R₃ is replaced by an oxygen atom.

37. (Withdrawn) A method according to claim 26, wherein R₁ is selected from alkyl and aminoalkyl.

38. (Withdrawn) A method according to claim 26, where R₁ is (S)-Methyl, (R)-Methyl or (S)-Propyl.

39. (Withdrawn) A method according to claim 26, where R₁ is (S)-Aminopropyl.

40. (Withdrawn) A method according to claim 26, wherein R₂ is selected from (R)-Aminomethyl-(imino)-propyl, (S)-Aminomethyl-(imino)-propyl or (S)-Methylthiomethyl.

41. (Withdrawn) A method according to claim 26, wherein R₃ is alkyl, aralkyl or substituted aralkyl.

42. (Withdrawn) A method according to claim 26, wherein R_3 is 3-bromophenethyl.
43. (Withdrawn) A method according to claim 26, wherein R_3 is 3,5 bis-(trifluoromethyl)phenethyl.
44. (Withdrawn) A method according to claim 26, wherein R_4 is aralkylamino.
45. (Withdrawn) A method according to claim 26, wherein R_4 is benzylamino
46. (Withdrawn) A method according to claim 26 (previously claim 24), wherein r_6 and r_7 with inclusion of n , is heterocyclyl.
47. (Withdrawn) A method according to claim 26 (previously claim 24), wherein $-nr_6r_7$ is isoindolinyl having the formula
48. (Withdrawn) A method according to claim 26, wherein R_5 is S.
49. (Withdrawn) A method according to claim 26, wherein R_5 is O.